

Reductions of (v_3) configurations

Marko Boben*

University of Ljubljana, FRI, IMFM
Slovenia

February 1, 2008

Abstract

Cubic bipartite graphs with girth at least 6 correspond to symmetric combinatorial (v_3) configurations. In 1887 V. Martinetti described a simple reduction method which enables one to reduce each combinatorial (v_3) configuration to one from the infinite set of so-called *irreducible* configurations. The aim of this paper is to show that a slightly extended set of reductions enables one to reduce each combinatorial (v_3) configuration either to the Fano configuration or to the Pappus configuration.

Key words: cubic graphs, incidence structures, configurations.

Math. Subj. Class. (2000): 05C62, 05C85, 68R05, 68R10.

1 Introduction

Although the results and methods presented in this article come out from the graph theory, it is maybe worth mentioning that the problem itself originates from incidence structures named combinatorial configurations and dates back to 1887 [9].

A *symmetric combinatorial configuration* of type (v_r) is an incidence structure of v points and v blocks (called lines) such that each line contains r points, each point belongs to r lines and each two different points are contained in at most one line.

One way to relate graphs and configurations is via their incidence graphs (or Levi graphs [11]). An *incidence graph* or *Levi graph* of the (v_r) configuration $\mathcal{C}$ is a bipartite r -regular graph on v *black* vertices representing the points of $\mathcal{C}$, v *white* vertices representing the lines of $\mathcal{C}$, and with an edge joining two vertices if and only if the corresponding point and line are incident in $\mathcal{C}$.

The property that two different points of a configuration are contained in at most one line implies that Levi graphs have girth (the length of the shortest cycle) at least 6. Conversely, each bipartite r -regular graph with girth at least 6 and with chosen black-white coloring determines precisely one (v_r) configuration. Note that if the coloring is not chosen, such graphs determine a pair

*Address: Marko Boben, Univerza v Ljubljani, Fakulteta za računalništvo in informatiko, Tržaška 25, 1000 Ljubljana, Slovenia; E-mail address: Marko.Boben@fri.uni-lj.si

of *dual configurations*, i.e. configurations with the role of their points and lines interchanged. More about graphs and configurations can be found in [1, 4, 7].

In 1887 V. Martinetti described the following construction method for symmetric (v_3) configurations [8, 9]. Suppose that in the given (v_3) configuration there exist two parallel (non-intersecting) lines $\{a_1, a_2, a_3\}$ and $\{b_1, b_2, b_3\}$ such that the points a_1 and b_1 are not on a common line. By removing these two lines, adding one new point c and three new lines $\{c, a_2, a_3\}$, $\{c, b_2, b_3\}$, $\{c, a_1, b_1\}$ we obtain a $((v+1)_3)$ configuration. It is not possible to obtain every (v_3) configuration from some $((v-1)_3)$ configuration by using this method. We call (v_3) configurations which can not be constructed in this way from a smaller one *A-irreducible configurations* and the others *A-reducible configurations*. We use the prefix A before the term (ir)reducible to distinguish this type of reduction from a more general one defined later in the paper.

Let us consider the Martinetti method on Levi graphs of (v_3) configurations. For the sake of simplicity we will use, as in [2], the notion (v_3) graph instead of Levi graph of (v_3) configuration, i.e. (v_3) graph is a bipartite cubic graph with girth ≥ 6 . We define A-reducible and A-irreducible (v_3) graphs corresponding to A-reducible and A-irreducible configurations respectively as follows. A (v_3) graph G is *A-reducible* if there exists an edge $uv \in E(G)$ such that $(G - \{u, v\}) + x_1y_1 + x_2y_2$ or $(G - \{u, v\}) + x_1y_2 + x_2y_1$ is again a (v_3) graph, where x_1, x_2 are the neighbors of u (different from v) and y_1, y_2 are the neighbors of v (different from u). Otherwise we call a (v_3) graph *A-irreducible*.

It turns out that there are four infinite families of connected A-irreducible graphs, which we present in the next section, plus the Pappus graph, i.e. the Levi graph of the Pappus configuration, see Figure 4. Note that Martinetti in his paper did not describe all families of irreducible graphs. This was not done until [2].

In this paper we show that a slightly more general definition of a (v_3) graph reduction (and hence (v_3) configuration reductions) leaves only two irreducible graphs: the Heawood graph, the Levi graph of the Fano configuration (projective plane of order 2), and the already mentioned Pappus graph.

An effort in this direction was made in 2000 by H. G. Carstens, T. Dinski, and E. Steffen [3], who gave a set of quite complicated reductions to show that they lead to the Heawood graph. But their claim is not entirely correct since it turns out that their reductions leave out the Desargues graph too (the Levi graph of the Desargues configuration) [12].

2 The main result

Given a bipartite cubic graph G , black vertex $u \in V(G)$, and white vertex $v \in V(G)$, let us say that we *reduce G by u and v* if we remove u and v and connect their neighbours in such way that G remains cubic and bipartite. If u and v are not connected then there are six ways to this, otherwise there are only two.

If it is possible to reduce a (v_3) graph G by some pair of its vertices such that the reduced graph is again a (v_3) graph, then we say that G is *B-reducible*; otherwise we call it *B-irreducible*. We notice that A-reduction is only a special case of the above definition where u and v are neighbouring vertices.

Our main result says that there are not many B-irreducible (v_3) graphs.

Theorem 1. *There are only two connected B-irreducible bipartite cubic graphs with girth at least 6, the Heawood graph and the Pappus graph.*

To prove this theorem we show that our reduction applies to all A-irreducible (v_3) graphs but the Heawood and the Pappus graph.

It is possible that the reduction of a connected graph makes it disconnected. So, the process of reducing a (v_3) graph may finally result in a graph whose connected components are only the Heawood and the Pappus graphs. But we may state a stronger result.

Theorem 2. *Every connected (v_3) graph can be reduced either to the Heawood graph or to the Pappus graph by the sequence of connected (v_3) graphs.*

Proof. Note that bipartite cubic graphs are 2-connected; this follows, for example, from [5, Corollary 6.4.6]. Hence the removal of two vertices produces at most three connected components in the general case and at most two in the case of A-reduction. If the addition of the necessary edges leaves the graph disconnected and this does not produce a 4-cycle, then there also exists a choice for the addition of the edges which gives a connected graph and does not produce a 4-cycle. (Such a choice exists since we add 3 edges in the general case and 2 in the case of A-reduction.) $\square$

2.1 A-irreducible graphs

Let us first list all A-irreducible graphs as they are presented in [2].

The first family of A-irreducible graphs. The first type of A-irreducible graphs can be described as follows. Let $T(n)$, $n \geq 1$, denote a graph on $20n$ vertices which is an union of n segments G_T shown in Figure 1 where the i -th segment ($i \geq 2$) and the $(i-1)$ -st segment are joined together by the edges $v_{i-1}^1 u_i^1$, $v_{i-1}^2 u_i^2$, $v_{i-1}^3 u_i^3$. We will use $T(n)$ in the following definitions. Let $T_1(n)$




Figure 1: The segment graph G_T defining the families $T_1(n)$, $T_2(n)$ and $T_3(n)$.

be the graph which is obtained from $T(n)$ by adding the edges $u_1^1 v_n^1$, $u_1^2 v_n^2$, and $u_1^3 v_n^3$. Let $T_2(n)$ be the graph obtained from $T(n)$ by adding the edges $u_1^3 v_n^1$, $u_1^2 v_n^2$, and $u_1^1 v_n^3$. And finally, let $T_3(n)$ be the graph obtained from $T(n)$ by adding the edges $u_1^1 v_n^3$, $u_1^2 v_n^1$, and $u_1^3 v_n^2$. See Figure 2.

It turns out that for each fixed $n \geq 1$ the graphs $T_1(n)$, $T_2(n)$, $T_3(n)$ are pairwise non-isomorphic A-irreducible (v_3) graphs.




Figure 2: The construction of the graphs $T_1(n)$ (a), $T_2(n)$ (b), and $T_3(n)$ (c) from $T(n)$ by adding three edges (shown thick) joining the last and the first segment.




Figure 3: Members of the second family of A-irreducible (v_3) graphs are graphs $D(n)$ with LCF notation $[5, -5]^n$, $n \geq 7$, [6].




Figure 4: The Pappus graph.

The second family of A-irreducible graphs. The second family of A-irreducible (v_3) graphs consists of graphs $D(n)$ described by LCF notation $[5, -5]^n$, $n \geq 7$, [6]. The notation $[5, -5]^n$ for G means that G is constructed from the cycle of length $2n$ to which chords joining the vertices at distance 5 and -5 (in reverse direction) are attached, see Figure 3. The smallest graph of this family, $D(7)$, is the Heawood graph. The members of this family are Levi graphs of the so-called cyclic (v_3) configurations with base line $\{0, 1, 3\}$.

The Pappus graph There exists another A-irreducible (v_3) graph which is not a member of the families described above. This is the Pappus graph, Levi graph of the Pappus configuration, see Figure 4.

The following theorem from [2] states that the graphs mentioned above are all A-irreducible graphs.

Theorem 3. *All connected A-irreducible (v_3) graphs are*

1. *graphs $D(n)$, i.e. graphs with LCF notation $[5, -5]^n$, $n \geq 7$,*
2. *the graphs $T_1(n)$, $T_2(n)$, $T_3(n)$, $n \geq 1$, and*
3. *the Pappus graph.*




Figure 5: Reduction of the first A-irreducible family.




Figure 6: Reduction of the second A-irreducible family.

2.2 Proof of the main Theorem

We conclude this section by the proof of Theorem 1.

Proof of Theorem 1. Because of Theorem 3 and the fact that A-reduction is a special case of B-reduction, we have to take into consideration only A-irreducible graphs. The reduction of any $T(n)$ graph goes as follows: remove w_1^1 and z_1^1 and add edges $u_1^1 y_1^3$, $u_1^3 y_1^1$, and $v_1^2 x_1^2$, see Figure 5.

The reduction of any graph $D(n)$, $n \geq 8$, goes as follows: remove u_2^1 and u_2^4 and add edges $v_1^1 u_2^3$, $u_2^2 v_2^2$, and $v_1^2 v_2^1$, see Figure 6.

In all cases it is easy to check that the girth of the reduced graph is still 6. In the case of the Heawood graph, $D(7)$, and the Pappus graph, any reduction leads to a graph with girth 4. $\square$

3 Conclusion

In the discussion above we have proved, in the language of configurations, that every combinatorial (v_3) configuration can be reduced either to the Fano configuration or to the Pappus configuration by a generalized Martinetti procedure. In this procedure we always remove one line and one point and correct the structure locally on the affected objects such that we again obtain a combinatorial configuration.

When speaking about combinatorial configurations we cannot avoid mentioning *geometric configurations*. A geometric (v_r) configuration is the structure of v points and v lines *in the plane* such that each point belongs to r lines and each line passes through r points. In the time when Martinetti wrote his article, the distinction between combinatorial and geometric configurations was not strong, if it existed at all. One indication that this was probably not the case is that he did not list the two smallest irreducible configurations (the Fano (7_3) configuration and the Möbius-Kantor (8_3) configuration) since neither of them is a geometric configuration.

Hence we may ask the following question. For which combinatorial configurations that can be realized as geometric configurations one can realize each of (at least one of) the reduced configurations? This is certainly not the case for the (9_3) configurations, but do there exist other ones?

Another question one may ask at the end is the following: which cubic bipartite graphs of girth at least g are irreducible if we do not allow the girth of the reduced graph to fall under g ? In Theorem 1 we have found graphs for $g = 6$, so the next step would be to find graphs for $g = 8$. Obviously, one graph is the smallest trivalent graph of girth 8 or *the Tutte 8-cage* (which is also bipartite) on 30 vertices. The second one is the unique (v_3) graph with girth 8 on 34 vertices (since there is no such graph on 32 vertices according to [1]). There are also larger graphs as the generalized Petersen graph $G(18, 5)$, while $G(26, 5)$ is reducible to a graph with girth 8.

More about bipartite graphs of large girth and their connection to configurations can be found in [10].

Acknowledgements

The research was supported in part by a grant from Ministrstvo za šolstvo, znanost in šport Republike Slovenije, grant P1-0294. The author would also like to thank to Tomo Pisanski for his encouragement and many fruitful suggestions during the preparation of this paper.

References

- [1] A. Betten, G. Brinkmann, T. Pisanski, *Counting symmetric configurations v_3* , Discrete Appl. Math. 99 (2000), 331–338.
- [2] M. Boben, *Irreducible (v_3) configurations and graphs*, Discrete Math., to appear.
- [3] H. G. Carstens, T. Dinski, E. Steffen, *Reduction of symmetric configurations n_3* , Discrete Appl. Math. 99 (2000), 401–411.
- [4] H. S. M. Coxeter, *Self-dual configurations and regular graphs*, Bull. Amer. Math. Soc. 56 (1950), 413–455.
- [5] R. Diestel, *Graph Theory*, Springer, New York, 1997.
- [6] R. Frucht, *Canonical representation of trivalent Hamiltonian graphs*, J. Graph Theory 1 (1977), 45–60.
- [7] H. Gropp, *Configurations and graphs*, Discrete Math. 111 (1993), 269–276.
- [8] B. Grünbaum, *Special Topics in Geometry – Configurations*, Math 553B, University of Washington, Seattle, 1999.
- [9] V. Martinetti, *Sulle configurazioni piane μ_3* , Annali di matematica pura ed applicata (2) 15 (1887-88), 1–26.
- [10] T. Pisanski, M. Boben, D. Marušič, A. Orbanić, A. Graovac, *The 10-cages and derived configurations*, Discrete Math. 275 (2004), 265–276.
- [11] T. Pisanski, M. Randić, *Bridges between Geometry and Graph Theory*, in *Geometry at Work*, C.A. Gorini, ed., MAA Notes 53 (2000).

- [12] E. Steffen, T. Pisanski, M. Boben, Reduction of symmetric configurations n_3 — Erratum, submitted.